

ASTR 1040: Stars & Galaxies



Gran Telescopio Canarias,
La Palma 10.4m

Prof. Juri Toomre TAs: Peri Johnson, Ryan Horton
Lecture 5 Tues 30 Jan 2018
zeus.colorado.edu/astr1040-toomre

Topics for Today

- What our atmosphere does to “light”
- Magic of “adaptive optics”
- Radio telescopes: many dishes make a big one (interferometry or “aperture synthesis”)
- Telescopes in space -- and why
- Our Nearest Star the Sun in overview
- Finish reading *Chap 14 (Our Star)* in detail
- Read *S4.1, S4.2 (Fundamental particles ..)*
- Observ Night report ...HW #1 returned graded

Some Events

- Lunar Eclipse (*Super Blood Blue Moon*)
tomorrow early: Earth shadow (umbra) touches 4:48am, reddish max at 6:30am, but sunrise ~7am (S-B observatory has telescope, binocs set up outside, but can see from anywhere)
- Go directly to *Fiske Planetarium* for this Thur class – please try to be there by 11am



REMINDER

Problems in Looking Through Our Atmosphere

- Many wavelengths are **absorbed** (just don't make it through to surface)
- Turbulence in atmosphere distorts light:
 - stars appear to “twinkle”
 - angular resolution is degraded
- Man-made light is reflected by air particles, yielding bright night sky
 - this is *light pollution*

Light Pollution



90% of Earth's population cannot see the Milky Way

How many light bulbs does it take to screw up an astronomer?

An immediately curable pollution: simply turn the lights off!

Stop "uplight", glare: wastes billions of \$\$ in energy, use "low pressure sodium"

Several famous observatories are now useless...

LA Basin View from Mt. Wilson Observatory, 1908 and 1998

Quest for Good Weather and Seeing

- **Mauna Kea, Big Island of Hawaii, 14,000' elevation, middle of the Pacific**
- **Dry, high, dark and isolated. Best on the planet?**

Telescope Sites

- The best sites are **high, dark and isolated.**
- Even in the best places, **atmospheric angular resolution is typically 0.3-0.5 arcsec at visible wavelengths**

Mauna Kea, Hawaii

Paranal, Chile

Kitt Peak, Arizona

Canary Islands

Reading Clicker Q

B

- Which wavelength regions CAN be studied with ground-based telescopes?
- A. All light with wavelengths longer than ultraviolet
- B. Radio, visible, and very limited portions of infrared and ultraviolet
- C. All light with wavelengths shorter than infrared
- D. Infrared, visible, and ultraviolet

Atmospheric Absorption of "Light"

- Earth's atmosphere absorbs most types of light (not entirely bad, or we would be dead!)
- Only **visible, radio, and some IR and UV** light get through to the ground


To observe other wavelengths, must put telescopes in space!

UV, X-rays and Gamma-rays

- These all have enough energy to ionize electrons out of atoms or break apart molecules
- Heavily absorbed by the atmosphere
- Space or high altitude (balloon, rocket) observatories are necessary

Why bother with "other light"?

- ... Many very hot objects shine brightest in such UV, X-ray and gamma-ray photons
- ... And cool star-forming regions are brightest in IR




"Hot new stuff" for Optical Observatories



Adaptive Optics (AO) – "de-twinkle" stars

- Wavefronts of star light are deformed by atmosphere
- Can distort shape of mirror (very fast) to correct for distortions by atmosphere – hot new technology



Adaptive Optics to the Rescue!


- Use a laser to create an artificial star and correct for the distortion caused by Earth's atmosphere
 - If you bounce the incoming light off a warped mirror (of exactly the right shape) the light comes off corrected
- It's like reversing the effect of a funhouse mirror



Adaptive Optics benefits


NEPTUNE

Without adaptive optics



May 24, 1999

With adaptive optics




June 27, 1999

Images from the Keck Observatory

Adaptive Optics wizardry

Galactic Center




Clicker Q - galaxy **B**

- In observing a distant galaxy, the H alpha spectral line of hydrogen (usually in the visible) is now in the IR portion of the spectrum. What can you conclude?
- A. Galaxy is made purely of hydrogen
- B. Galaxy is moving away from us
- C. Galaxy is moving towards us
- D. Galaxy has very weak gravity

So what gets through our atmosphere?

- **RADIO WAVES:** most get through
 - Thus radio telescopes are built on the ground
- Weather is not an issue
 - Radio waves come right through the clouds
- But poorer angular resolution
 - Why?
 - **VERY long wavelengths!**




300 feet

Green Bank Telescope, West Virginia



Arecibo, Puerto Rico


1000 feet!




Interferometry

- Join **multiple telescopes together** to simulate **one large telescope**.
- **Very Large Array (VLA) in New Mexico** has 27 dishes (each 25 m) across in a 40 km valley
- **Very Large Baseline Array (VLBA)** is an array of ten telescopes around the hemisphere
 - Resolutions as small as 0.001 arcseconds for radio light
- The **twin Keck telescopes** can also be an infrared interferometer.


Very Large Array (VLA)
New Mexico




Very Large Baseline Array (VLBA)



Keck Telescopes
Hawaii






VLA – “Large Aperture Synthesis”

Radio Interferometry – many small look big!

- Two (or more) radio dishes observe the same object
- Signals from each “interfere” with each other
- Can construct image whose angular resolution is like that from a huge dish!



Very Large Array (VLA) NM

27 25m antennas
36 km baseline

VLBA in Owens Valley CA

continental baselines 10 25m dishes

ALMA: Atacama Large mm-submm (M) Array
(0.3-3.6 mm) 84-950 GHz

Very high desert Chile (altiplano 5000 m !)

One of 50+
12m diam antennas
15km max separation
(0.005" best resolution)
but typical res 0.1"

ALMA

Palomar Observatory 5,600 FT
Kitt Peak Observatory 6,900 FT
Very Large Array 7,000 FT
Keck Observatory 14,000 FT
ALMA Observatory 16,500 FT

CCAT – 25 m “wide-angle” sub-mm telescope CU is partner

Located at 5600m altitude (above ALMA)

Infrared Telescopes

- **INFRARED** can be absorbed by molecules (mostly H₂O) in the Earth's atmosphere.
- Two recent solutions:
 - Fly above the clouds!
 - Go where there is no water!

For other wavelengths we have to get above the atmosphere

- UV, X-rays, Gamma Rays
- **Methods:** balloons, rockets, Space Shuttle, satellites



NASA's Great Observatories

- Compton Gamma Ray Observatory
- Spitzer Space Telescope Infrared
- Hubble Space Telescope UV/Visible
- Chandra X-Ray Observatory




How do you point a space telescope in orbit ?

1. Squirt from jets to change direction (hydrazine)
2. Torque by electric currents in big coils while flying through Earth's magnetic field
3. Torque by electric motors spinning up or down "reaction wheels"

ANGULAR MOMENTUM DEMONSTRATION

Hubble Space Telescope: NASA's most famous observatory

- Launched in 1990
 - Error in mirror made blurry images
- Corrective optics installed in 1993 (Ball Aerospace here in Boulder)
- Small (only 2.5 meters) but diffraction-limited
- Low orbit accessible by Shuttle, refurbishing missions mean long lifetime (1990 to 2014+)
- \$5 billion over 20 years = 10-100 times more costly than ground-based telescope




Very sharp images from Hubble ... and much more





"Nonvisible" Light – X-ray, UV, IR, Radio

- Most light is invisible to human eye
- Special detectors can record such light
- Digital images built using false-color coding



Chandra X-ray image of center of our Milky Way Galaxy


SPITZER Infrared Telescope

- Launched August 2003
- **Trails behind Earth to get away from Earth's thermal spectrum**
- **0.85m aperture , T ~ 5.5 K**
- Cooled with liquid helium, had 2-5 years worth, now used up (warmer phase)

X-Ray Telescopes – do it their own way!

- X-ray photons can pass right through a mirror
- Such photons can only be **reflected at shallow angles**, like “skimming stones” off water surface



Chandra X-Ray Observatory

mirrors of first reflection mirrors of second reflection

X rays

focal point

(a) X rays

(b) X rays

(c) doubly reflected X rays


second set of four nested mirrors

first set of four nested mirrors

focal plane

10 meters

Bigger view of Chandra's X-ray Imaging



X rays

X rays

The first set of nested mirrors deflects X rays to the second set . . .

10 meters

focus


“grazing reflections”

. . . and the second set deflects X rays a second time to the focus point.

Chandra X-ray Observatory (at L₂ Lagrangian)



Multi-energy X-ray picture



lowest-energy X rays (red)

medium-energy X rays (green)

highest-energy X rays (blue)

Chandra: Supernova Remnant SNR: N132D

Clicker

What does the technique of interferometry allow?

- ... to make astronomical observations without interference from light pollution
- ... the same telescope to make images with both radio waves and visible light
- ... to determine the chemical composition of stars
- ... multiple telescopes to obtain the angular resolution better than the individual telescopes
- ... multiple telescopes to obtain a total light-collecting area larger than the individual telescope


Instruments in the Focal Plane

How astronomers use light collected by a telescope:

1. **Imaging**
 - use camera to take pictures (images)
 - photometry: measure amount and color (with filters) of light from object
2. **Spectroscopy**
 - use spectrograph to separate light in detail into its different wavelengths (colors)
3. **Timing**
 - measure how amount of light changes with time (sometimes in a fraction of a second)


Imaging (Digital with CCDs)

- **Filters** are placed in front of camera to allow only certain colors to be imaged
- Single color **images** are superimposed to form “true color” images.




Spectroscopy – analyzing the light


- Spectrograph reflects light off a **grating**: finely ruled, smooth surface
- Light (by interference) disperses into colors
- This **spectrum** is recorded by digital CCD detector



Spectral resolution is vital but also “costly in photons”



Light curves: Studying changes with time



Variable star MIRA: period ~331 days